


AD-A173 506 CHIKUNGUNYA FEVER AMONG US PEACE CORPS VOLUNTEERS - 1/1
REPUBLIC OF THE PHILIPPINES(U) NAVAL MEDICAL RESEARCH
UNIT NO 2 MANILA (PHILIPPINES) C HAYES ET AL
UNCLASSIFIED 12 SEP 86 NANRU-2-TR-1017 F/G 6/13 NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A173 506


2

CHIKUNGUNYA FEVER AMONG U.S. PEACE CORPS VOLUNTEERS -
REPUBLIC OF THE PHILIPPINES

C.G. Hayes, T. O'Rourke and A. Sarr.

REPORT NO.

TR - 1017



DTIC
ELECTE
OCT 24 1986
S B

DTIC FILE COPY

**UNITED STATES NAVAL
MEDICAL RESEARCH UNIT NO. TWO**

APO SAN FRANCISCO, CALIFORNIA 96528

NAVAL MEDICAL RESEARCH AND DEVELOPMENT COMMAND
BETHESDA, MARYLAND

86 10 16 183

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE

ADA173506

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION Unclassified		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION / AVAILABILITY OF REPORT Distribution of this document is unlimited.	
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE			
4. PERFORMING ORGANIZATION REPORT NUMBER(S) NAMRU-2-TR-1017		5. MONITORING ORGANIZATION REPORT NUMBER(S)	
6a. NAME OF PERFORMING ORGANIZATION U.S. Naval Medical Research Unit No. 2	6b. OFFICE SYMBOL (If applicable) NAMRU-2	7a. NAME OF MONITORING ORGANIZATION	
6c. ADDRESS (City, State, and ZIP Code) APO San Francisco, California 96528		7b. ADDRESS (City, State, and ZIP Code)	
8a. NAME OF FUNDING / SPONSORING ORGANIZATION Naval Medical Research & Development Command	8b. OFFICE SYMBOL (If applicable) NMRDC	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER	
8c. ADDRESS (City, State, and ZIP Code) Bethesda, Maryland 20814		10. SOURCE OF FUNDING NUMBERS	
		PROGRAM ELEMENT NO. 61102A	PROJECT NO. 3M161102FSLD TASK NO. AF WORK UNIT ACCESSION NO. 429
11. TITLE (Include Security Classification) (U) Chikungunya fever among U.S. Peace Corps Volunteers - Republic of the Philippines			
12. PERSONAL AUTHOR(S) Curtis Hayes, Thomas O'Rourke, A. Sarr Center for Infectious Diseases, CDC.			
13a. TYPE OF REPORT Technical Report	13b. TIME COVERED FROM 1985 TO 1986	14. DATE OF REPORT (Year, Month, Day) 1985 June to 1986 Jan.	15. PAGE COUNT
16. SUPPLEMENTARY NOTATION Published in the MMWR, 35(36):573-574, Sept. 12, 1986. Prepared in cooperation with the U.S. Peace Corps, Philippines; and Center for Infect. Dis.			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	
		chikungunya fever Philippines	
		U.S. Peace Corps	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Three cases of chikungunya fever were diagnosed recently among U.S. Peace Corps volunteers stationed in the Republic of the Philippines. These are the first cases reported from the Philippines since 1968. All three cases were diagnosed using an indirect antibody-capture enzyme-linked immunosorbent assay (ELISA). The cases were diagnosed as part of a collaborative long-term infectious disease study by the U.S. Naval Medical Research Unit No. 2 and the U.S. Peace Corps in the Philippines.			
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED / UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL RAM I: M. Groves		22b. TELEPHONE (Include Area Code) 301/663-6576	22c. OFFICE SYMBOL NMRDC

DD FORM 1473, 84 MAR

83 APR edition may be used until exhausted.
All other editions are obsolete.

SECURITY CLASSIFICATION OF THIS PAGE

Reproduced at Government Expense

Unclassified

ADMINISTRATIVE INFORMATION

C.G. HAYES, Ph.D.
Acting Scientific Director

This study was supported through funds provided by the U.S. Naval Medical Research and Development Command for Work Unit No. 34161102 BS10 AF429.

Distribution of this document is unlimited.

L. W. LAUGHLIN
CAPT MC USN

Accession	
NTIS	
DUC	
Under	
JUL	
GPO	
By	
Distribution	
Availability	
Dist	Avail
A-1 24	Spec



Stock #717-016-00100-5
Price: \$1.00

International Notes

**Chikungunya Fever among U.S. Peace Corps Volunteers —
Republic of the Philippines**

Three cases of chikungunya fever were diagnosed recently among U.S. Peace Corps volunteers stationed in the Republic of the Philippines. These are the first cases reported from the Philippines since 1968. The cases were diagnosed as part of a collaborative long-term infectious disease study by the U.S. Naval Medical Research Unit No. 2 and the U.S. Peace Corps in the Philippines.

The first patient, a 27-year-old male, was stationed in Mindanao, one of the southernmost islands of the Philippine archipelago. His illness occurred in June 1985 but was not diagnosed until October 1985. The second case occurred mid-November 1985 in a 31-year-old female stationed on Cebu, one of the islands that forms the central portion of the archipelago. The third case occurred in January 1986 in a 23-year-old female stationed on the island of Masbate, also in the central part of the country but north of Cebu.

Chikungunya Fever — Continued

All three cases were diagnosed using an IgM antibody-capture enzyme-linked immunosorbent assay (ELISA). Chikungunya virus was isolated from a blood sample obtained from one of the patients 3 days after onset of fever. The clinical presentations were typical of chikungunya fever and included acute onset of high fever, severe joint pain, and skin rash. The illnesses persisted 3-7 days, and all patients recovered uneventfully.

Chikungunya virus may have been introduced into the southern Philippines from Indonesia. The first case detected in the Philippines occurred in an area of Mindanao, Davao del Sur, which is approximately 200 km north of central Indonesia; the area is reportedly frequently visited by Indonesian traders and fishermen.

The full extent of epidemic activity in the Philippines is not known. One of the Peace Corps volunteer patients reported many cases of a chikungunya-like illness (locally termed "Chinese fever" or "Asian flu") in the Filipino population of the village where she lived. The potential for widespread activity is present because of the large population of susceptibles and the abundance of the vector, *Aedes aegypti*, particularly in large urban areas like Manila.

Reported by CG Hayes, PhD, T O'Rourke, MD, U.S. Naval Medical Research Unit No. 2; A Sarr, US Peace Corps, Philippines, Div of Vector-Borne Viral Diseases, Center for Infectious Diseases, CDC.

Editorial Note: The word "chikungunya" is Swahili for "that which bends up," in reference to the stooped posture of patients afflicted with the severe joint pain associated with this disease. The disease was first recognized in epidemic form in East Africa in 1952-1953 (1). The etiologic agent, chikungunya virus, is arthropod-borne and has been placed in the family *Togaviridae*, genus *Alphavirus* (2). Human infections are acquired by the bite of infected *A. aegypti* mosquitoes, and epidemics are sustained by human-mosquito-human transmission. The epidemic cycle is thus similar to those of dengue and urban yellow fever.

Since 1954, the virus has been implicated as the cause of epidemics in Asian countries including the Philippines, Thailand, Cambodia, Vietnam, India, Burma, and Sri Lanka. Epidemics of chikungunya were documented in the Philippines in 1954, 1956, and 1968 (3-4). Serosurveys suggest that virus activity occurred in the central and southern part of the archipelago (5). Chikungunya virus has been isolated from humans and mosquitoes in eastern, southern, western, and central Africa and in southeastern Asia, where it has been responsible for illnesses in hundreds to thousands of individuals.

Chikungunya fever is characterized by sudden onset, chills and fever, headache, nausea, vomiting, arthralgia, and rash. In contrast to dengue, chikungunya is characterized by a briefer febrile episode, by persistent arthralgia in some cases, and by the absence of fatalities. However, similarities between clinical appearances of the two diseases probably account for misclassification and some underreporting of chikungunya fever in areas with endemic dengue; therefore, laboratory confirmation of reported cases is important. The IgM-capture ELISA used for diagnosis of these patients is especially useful in this regard (6).

References

1. Robinson MC. An epidemic of virus disease in Southern Province, Tanganyika Territory, in 1952-1953. I. Clinical features. *Trans Royal Society Trop Med Hyg* 1955;49:28-32.
2. Calisher CH, Shope RE, Brandt W, et al. Proposed antigenic classification of registered arboviruses. I. *Togaviridae*, *Alphavirus*. *Intervirology* 1980;14:229-32.
3. Campos LE, San Juan A, Cenabre LC, Almagro EF. Isolation of chikungunya virus in the Philippines. *Acta Med Philippina* 1969;5:152-5.
4. Macasaet FF. Further observation on chikungunya fever. *J Philippines Med Assoc* 1970;46:235-42.
5. Tesh RB, Gadjusek DC, Garruto RM, et al. The distribution and prevalence of group A arbovirus neutralizing antibodies among human populations in Southeast Asia and the Pacific Islands. *Am J Trop Med Hyg* 1975;24:664-75.
6. Calisher CH, El-Kafrawi AO, Al-Deen Mahmud MI, et al. Complex-specific immunoglobulin M antibody patterns in humans infected with alphaviruses. *J Clin Microbiol* 1986;23:155-9.

END

12-86

DTIC